

Expanding Universe : Thermodynamical Aspects From Different Models

Sridip Pal ^{*}1

Ritabrata Biswas [†]2

** Department of Physical Sciences, Indian Institute of Science Education and Research, Mohanpur - 741252, Nadia, West Bengal, India*

† Department of Physics, Indian Institute of Science, Bangalore-560012, India.

Abstract

The pivotal point of the paper is to discuss the behavior of temperature, pressure, energy density as a function of volume along with determination of caloric EoS from following two model: $w(z) = w_0 + w_1 \ln(1+z)$ & $w(z) = -1 + \frac{(1+z)}{3} \frac{A_1 + 2A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2}$. The time scale of instability for this two models is discussed. In the paper we then generalize our result and arrive at general expression for energy density irrespective of the model. The thermodynamical stability for both of the model and the general case is discussed from this viewpoint. We also arrive at a condition on the limiting behavior of thermodynamic parameter to validate the third law of thermodynamics and interpret the general mathematical expression of integration constant U_0 (what we get while integrating energy conservation equation) physically relating it to number of micro states. The constraint on the allowed values of the parameters of the models is discussed which ascertains stability of universe. The validity of thermodynamical laws within apparent and event horizon is discussed.

PACS No : 05.70.-a (Thermodynamics), 95.30.Tg (Thermodynamic processes, conduction, convection, equations of state)

¹sridippaliiser@gmail.com

²biswas.ritabrata@gmail.com, ritabrata@physics.iisc.ernet.in

Contents

1	Introduction	2
2	Study of Universe Treating it as an Thermodynamical System	4
2.1	Thermodynamics of Fluid with $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$	5
2.2	Thermodynamics of Fluid with $\omega(z) = -1 + \frac{1+z}{3} \frac{A_1+2A_2(1+z)}{A_0+2A_1(1+z)+A_2(1+z)^2}$	6
3	Derivation of Thermal EoS & Study of Stability	7
3.1	Fluid with $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$	7
3.2	Fluid with $\omega(z) = -1 + \frac{1+z}{3} \frac{A_1+2A_2(1+z)}{A_0+2A_1(1+z)+A_2(1+z)^2}$	8
4	A General Derivation : Independent of Model	10
5	Instability : Time scale for Onset of Instabilities	11
6	On the Validity of Thermodynamical Laws	12
6.1	Validity on Apparent Horizon	12
6.2	Invalidity of Thermodynamic laws on Event Horizon	13
7	Brief Summary	14

1 Introduction

To explain the cosmic acceleration predicted from the Ia type supernova observations (Perlmutter, S. et al. 1999; Riess, A. G. et al. 1998) one popular way out is to modify the stress energy tensor part, i.e., the right hand side of the Einstein's field equation. Existence of some unknown matter termed as Dark Energy(DE hereafter) is been assumed (Riess, A. G. et al 2004; Perlmutter, S. et al 1998; Garnavich, P. M. et al 1998; Bachall, N. A. et. al. 1999; Copeland, E. J. et. al. 2006) which violates the strong energy condition. The simplest candidate of DE is a tiny positive cosmological constant(Λ) which obeys the equation of state (EoS hereafter), $w = -1$. But due to low energy scale than the normal scale for constant Λ , the dynamical Λ was introduced (Caldwell, R. R. et. al. 1998). Again at very early stage of universe the energy scale for varying Λ is not sufficient. So to avoid this problem, known as cosmic coincidence (Steinhardt, P. J. et. al. 1999), a new field, called tracker field (Zlatev, I. et. al. 1999) was prescribed. In similar way there are many models (Sahni, V. and Starobinsky, A. A. 2003) in Einstein gravity to best fit the data. Yet its require some modifications. From this point of view some alternative models are evolved. Most of the DE models involve one or more scalar fields with various actions and with or without a scalar field potential (Maor, I. et. al. 1998). Now, as the observational data permits us to have a rather time varying EoS, there are a bunch of models characterized by different scalar fields such as a slowly rolling scalar field (Quintessence) ($-1 < \omega < -1/3$, $\omega(= p/\rho)$, being the EoS parameter) (Caldwell, R. R. et. al. 1998), k-essence (Armendariz - Picon et. al. 2000), tachyon (Sen, A. 2002), phantom ($\omega < -1$) (Caldwell, R. R. 2002), ghost condensate (Arkani-Hamed, N. et. al. 2004; Piazza, F and Tsujikawa, S. 2004), quintom (Feng, B. et. al. 2005), Chaplygin gas models (Kamenshchik, A. Y. et. al. 2001) etc. Some recent reviews on DE models are described in the ref. (Copeland, E.J. et. al. 2006; Li, M. et. al.2011).

While explaining evolution of the universe, various DE models have been proposed, all of which must be constrained by astronomical observations. In all the models, the EoS parameter ω plays a key role and can reveal the nature of DE which accelerates the universe. Different equations of state lead to different dynamical changes and may influence the evolution of the universe. The EoS parameter ω and its time derivative with respect to Hubble time are currently constrained by the distance measurements of the type Ia supernova and the current observational data constrain the range of equation of state as $-1.38 < \omega < -0.82$ (Melchiorri, A. et. al. 2003). Recently, the combination of WMAP and Supernova Legacy Survey data shows a significant constraint on the EoS $\omega = -0.97^{+0.07}_{-0.09}$ for the DE, in a flat universe (Seljak, U. et. al. 2006). Recently some parametrization for the variation of EoS parameters $\omega(z)$ have been proposed describing the DE component.

1. $\omega(z) = \omega_0 + \omega_1 z$ (Cooray, A. R. and Huterer, D. 1999). Here $w_0 = -1/3$ and $w_1 = -0.9$ with $z < 1$. This grows increasingly unsuitable for $z > 1$. So the following model has been proposed. We will call this parametrization as '*linear parametrization*'.

2. $\omega(z) = \omega_0 + \omega_1 \frac{z}{1+z}$. This ansatz was first discussed by Chevallier and Polarski (Chevallier, M. and Polarski, D. 2001) and later studied more elaborately by Linder (Linder, E. V. 2003). The best fit values for this model while fitting with the SNIa gold dataset are $\omega_0 = -1.58$ and $\omega_1 = 3.29$. We will call this parametrization as '*CPL(Chevallier – Polarski – Linder) parametrization*'.

3. $\omega(z) = \omega_0 + \omega_1 \frac{z}{(1+z)^2}$ (Jassal, H. K. et. al. 2005). A fairly rapid evolution of this EoS allowed so that $\omega(z) \geq -1/2$ at $z > 0.5$ is consistent with the supernovae observation. We will call this parametrization as '*JBP(Jassal – Bagla – Padmanabhan) parametrization*'.

4. $\omega(z) = -1 + \frac{(1+z)}{3} \frac{A_1 + 2A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2}$ (Alam, U. et. al. 2004a, 2004b). This ansatz is exactly the cosmological constant $\omega = -1$ for $A_1 = A_2 = 0$ and DE models with $w = -2/3$ for $A_0 = A_2 = 0$ and $\omega = -1/3$ for $A_0 = A_1 = 0$. It has also been found to give excellent results for DE models in which the equation of state varies with time including quintessence, Chaplygin gas, etc. The best fit values of A_1 and A_2 are $A_1 = -4.16$ and $A_2 = 1.67$ for the SNIa Gold dataset. We will call this parametrization as '*ASSS(Alam – Sahni – Saini – Starobinski) parametrization*'.

5. $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$ (Efsthathiou, G. 1999). This evolution form of EoS is valid for $z < 4$. ω_1 is a small number which can be determined by the observations. The minimum value of ω_1 is approximately -0.14 and $\omega_0 \geq -1$. We will call it '*Log parametrization*'.

Thermodynamics of DE universe filled up with the fluids with linear and JBP parametrization had been studied before (Mazumder, N. et. al. 2010). Thermodynamics with CPL was studied by Xing, L. et. al. (2011). In this paper we are going to study DE universe filled with fluid Log parametrization and ASSS parametrization respectively which have not been studied before so extensively.

As we went through the literature we came to know that apart from the cosmological constant ($\omega = -1$), the sinehyperbolic scalar field potential (Sahni, V. and Starobinsky, A.A 2000; Urena-Lopez, L.A. and Matos, T. 2000; Sahni, V. et al. 2003) and the topological defect models, there is no DE model with constant ω consistent with observation (Alam, U. et. al. 2004a). Almost every model like Quintessence, Chaplygin gas (Setare, M.R. and Bouhmadi-Lopez, M.R. 2007, Setare, M.R. 2007b) etc. depicts significant evolution in $\omega(z)$ over sufficient time scale.

The way towards a meaningful reconstruction of $\omega(z)$ depends on inventing an efficient fitting function for either $d_L(z)$ or $H(z)$, where $d_L(z)$ is luminosity distance and $H(z)$ is Hubble parameter. The parameters of this fitting function are determined by matching to Supernova observations. Now, one can manage to reconstruct the functional form of $\omega(z)$ by taking an ansatz of the luminosity distance or $H(z)$ and then by comparing it to supernova observations.

This model is based on the following ansatz:

$$H(x) = H_0 (\Omega_m x^3 + A_0 + A_1 x + A_2 x^2)^{\frac{1}{2}}. \quad (1)$$

Which can be equivalently written as following:

$$\rho_{DE} = \rho_{0c} (A_0 + A_1 x + A_2 x^2 + A_3 x^3) \quad (2)$$

where $x = (1+z)$, ρ_{0c} is the critical density at present time and Ω_m dimensionless density of dark energy.

If we carefully note we can see the ansatz in terms of energy density is actually a truncated Taylor expansion where it is exploited that any well behaved function can be well approximated by Taylor expansion within a range. So the model is valid for $z \leq \text{few}$ (Alam U et. al. 2004a).

The model yields excellent results among DE models in which the EoS varies with time including quintessence, Chaplygin gas etc. (Sahni, V. et al. 2003; Alam, U. et al. 2003). Alam, U et al. 2004a (In figure 1 of their paper) have shown the accuracy of the ansatz when it is applied to several other DE models like tracker quintessence, the Chaplygin gas and super-gravity (SUGRA) models. They also plotted the deviation of $\log(d_L H_0)$ (which is the measured quantity for SNe) obtained via the ansatz from the actual model values. Clearly the ansatz comes out to be of excellent agreement well over a significant red-shift range for $\Omega_{0m} = 0.3$. The ansatz agrees

with these models of dark energy with less than 0.5% errors in the redshift range where we do have SNe data available (Alam, U. et. al. 2004a)

The analysis of type Ia supernova data involving the priors with most frequently used condition as $\omega = \text{constant}$ and $\omega \geq -1$ leads to confinement of DE to within a narrow class of models. Moreover, when we impose such priors on the cosmic equation of state it can result to a complete misrepresentations of reality as shown in (Maor, et al. 2002).

Recently, evolution of EoS parameter with red shifts been investigated (Vazquez, J. A. et. al. 2012) by performing a Bayesian analysis of current cosmological observations. Vazquez, J. A. et. al. have shown if they calculate the Bayes' factor then most of the data catalogues supports the fact that CPL, JBP are significantly disfavored with respect to simple Λ CDM model ($\omega = -1$). Now, linear parametrization is not to be considered in such a case as the parametrization blows with high z . Calculation of Bayes' factor for ASSS and Λ CDM using codes like CAMB etc are bit cumbersome. But from intuition we can see -1 term is already there in ASSS parametrization. Extra additional z dependent term may make it different enough from the Λ CDM model. Even theoretically, ASSS is efficient enough to explain different phases of universe. These observational supports substantiate our interest of working with ASSS model.

Linear, CPL, JBP or ASSS - all the parametrizations can be treated as kind of combinations of polynomials with $(1+z)$ as polynomial parameter. These modes are efficient for early epochs whereas they are not that effective for low redshifts. While parametrizing, the fact that at low redshifts the magnitude-redshift relation is degenerate for the models having same deceleration parameter, should be taken into account. Besides keeping consistency of SN constraints on dimensionless DE density with those derived from CAMB measurements is another important factor. These constraints along with many others had given birth of Log parametrization. This is somehow not following the common pathway of considering DE EoS. These properties did motivate us to take Log parametrization for the study of thermodynamics of universe.

We will study the universe from thermodynamical aspect for both type of fluids separately in Sec. 2. In Sec. 3 we will seek the thermal EoS for both type of fluid separately along with study on thermodynamical stability. The Internal Energy, thermodynamic pressure, entropy, temperature as a function of volume have been determined in this section. The Sec. 4 contains a general derivation of thermal EoS for any fluid obeying $p = \omega(z)\rho$ and a general discussion on stability criterion. We have tried to give a physical interpretation of the generality of thermal EoS and the expression for energy density as a function of temperature and volume in this section. Study of instable cases, onset of instability have been discussed in Sec. 5. The Sec. 6 proves the validity of laws of thermodynamics on the apparent horizon and invalidity of the same on the event horizon which agrees with (Mazumder, N. et. al. 2010; Xing, L. et. al. 2011). The paper ends with a brief concluding remark in Sec. 7.

2 Study of Universe Treating it as an Thermodynamical System

Let us consider an Universe filled with a perfect fluid having volume V and ρ , p , T and S are respectively the energy density, thermodynamical pressure, temperature and entropy of the system. From the first law of thermodynamics (Myung, Y. S. 2011) we have

$$TdS = d(\rho V) + pdV = d[(\rho + p)V] - Vdp \quad (3)$$

We have the following integrability condition (Myung, Y. S. 2011), i.e.,

$$\frac{\partial^2 S}{\partial T \partial V} = \frac{\partial^2 S}{\partial V \partial T}$$

which yields (Gong, Y. et. al. 2007)

$$\frac{dp}{\rho + p} = \frac{dT}{T}. \quad (4)$$

So combining equations (3) and (4) and integrating we obtain the following:

$$S = \frac{(\rho + p)V}{T}, \quad (5)$$

where we have dropped an additive constant devoid of physical significance.

We assume our universe to be homogeneous and isotropic FRW space-time with following line element

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right], \quad (6)$$

where k , the curvature scalar having values 0, ± 1 for flat, closed and open universe respectively. The Friedmann equations and the energy conservation equation are

$$H^2 = \frac{8\pi G\rho}{3} - \frac{k}{a^2}, \quad (7)$$

$$\dot{H} = -4\pi G(\rho + p) + \frac{k}{a^2} \quad (8)$$

and

$$\dot{\rho} + 3H(\rho + p) = 0, \quad (9)$$

where the Hubble parameter is given by $H = \frac{\dot{a}}{a}$.

2.1 Thermodynamics of Fluid with $\omega(z) = \omega_0 + \omega_1 \ln(1 + z)$

Using this $\omega(z)$, integrating the energy conservation equation (9) we would get the expression of energy density as a function of redshift as follows :

$$\rho = \rho_0(1 + z)^{3\{1 + \omega_0 + \frac{\omega_1}{2} \ln(1+z)\}}, \quad (10)$$

where ρ_0 is integration constant denoting the present time ($z = 0$) density of our universe.

1. From the graph of energy density vs redshift, we observe that ρ initially has a very high value. With expansion of the universe, ω decreases, hence ρ falls to ρ_0 as z becomes 0 from 2.
2. The graph of Pressure vs Redshift reveals that the pressure remains positive for allowed values of ω_0 and ω_1 in the range $z \in (0, 2)$. It initially starts with high value and then decreases down to present value as z goes from 2 to 0.

The integration of the integrability condition (4) gives expression for temperature as

$$T = T_0(1 + \omega)(1 + z)^{3\{\omega_0 + \frac{\omega_1}{2} \ln(1+z)\}}, \quad (11)$$

where T_0 is the integration constant (Note $T_0(1 + \omega_0)$ is the present time temperature).

- The graph between Temperature and Redshift reveals that Temperature is also dropping as we traverse from past to present, i.e, as z varies from 2 to 0.


Using the last two expressions and plugging in the expression for entropy in the last section, we get the following expression for S :

$$S = \frac{\rho_0}{T_0}. \quad (12)$$

The heat capacity and square of the sound velocity are given by:

$$c_V(z) = V \frac{\partial \rho}{\partial T} = \frac{3S(1 + \omega)}{\omega_1 + 3\omega(1 + \omega)}. \quad (13)$$

Note when $\omega < -1$ we have $c_V < 0$ i.e universe is in a unstable phase. Hence the stability condition demands: $\omega > -1$.



The expression for the sound's speed :

$$v_s^2(z) = \frac{\partial p}{\partial \rho} = \omega + \frac{\omega_1}{3(1+\omega)}. \quad (14)$$

Now demanding $v_s^2 < 1$ we get the following condition:

$$\{\omega_0 + \omega_1 \ln(1+z)\}^2 \leq \frac{3-\omega_1}{3}, \quad (15)$$

along with the constraint $\omega_1 \leq 3$. z has a minimum value of 0, hence we can write:

$$3\omega_0^2 + \omega_1 \leq 3. \quad (16)$$

2.2 Thermodynamics of Fluid with $\omega(z) = -1 + \frac{1+z}{3} \frac{A_1+2A_2(1+z)}{A_0+2A_1(1+z)+A_2(1+z)^2}$

Using this $\omega(z)$, integrating the energy conservation equation we would get the following:

$$\rho = \rho_0 M \exp \left\{ -\frac{A_1}{\sqrt{A_0 A_2 - A_1^2}} \arctan \left(\frac{A_0 + A_2(1+z)}{\sqrt{A_0 A_2 - A_1^2}} \right) \right\}, \quad (17)$$

where ρ_0 is integration constant which we can get putting $z = 0$, and $M = A_0 + 2A_1(1+z) + A_2(1+z)^2$.

On the other hand integrating the integrability condition, we get the expression for temperature:

$$T = T_0(1+\omega)(1+z)^{-3} M \exp \left\{ -\frac{A_1}{\sqrt{A_0 A_2 - A_1^2}} \arctan \left(\frac{A_0 + A_2(1+z)}{\sqrt{A_0 A_2 - A_1^2}} \right) \right\}, \quad (18)$$

where T_0 is the integration constant.

Using the last two expressions and plugging in the expression for entropy in the last section, we get the following expression for S :

$$S = \frac{\rho_0}{T_0}. \quad (19)$$

The heat capacity and square of the sound velocity are given by:

$$c_V(z) = V \frac{\partial \rho}{\partial T} = S \left\{ \frac{1}{1+\omega} + \frac{\frac{3}{(1+\omega)(1+z)} - \frac{1}{(1+\omega)^2} \frac{\partial \omega}{\partial z}}{\frac{1}{1+\omega} \frac{\partial \omega}{\partial z} + \frac{3\omega}{1+z}} \right\}. \quad (20)$$

Upon simplification we get:

$$c_V = \frac{3S(2+\omega)}{3(1+\omega)^2 + (1+z) \frac{\partial \omega}{\partial z}} \quad \text{and} \quad (21)$$

$$v_s^2(z) = \frac{\partial p}{\partial \rho} = \omega + \frac{\partial \omega}{\partial z} \left\{ \frac{A_1 + A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2} \right\}^{-1} \quad (22)$$

3 Derivation of Thermal EoS & Study of Stability

In this approach it is very convinient to consider the fluid obeying adiabatic EoS $p = \omega(z)\rho$ with constant particle number N as a thermodynamical system. Without any loss of generality we can assume the internal energy (U) and the pressure (p) as functions of entropy (S) and volume (V). So we can structurize our density and pressure and also the concerend differential equation as (Landau, L. D. and Lifschitz, E. M. 1984)

$$\left. \begin{aligned} \rho &= \frac{U}{V}, \\ p &= -\left(\frac{\partial U}{\partial V}\right)_S \quad \text{and} \\ \frac{dU}{dV} + \omega(z)\frac{U}{V} &= 0. \end{aligned} \right\} \quad (23)$$

3.1 Fluid with $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$

In (23), use of $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$, yields the following:

$$U = U_0 V^{(\frac{\omega_1}{6} \ln V)} \exp(-\omega_0 V), \quad (24)$$

where U_0 is integration constant (It can be function of S and ω_0 should be greater than 0 for U not to diverge). Hence the energy density becomes

$$\rho = U_0 V^{(\frac{\omega_1}{6} \ln V - 1)} \exp(-\omega_0 V), \quad (25)$$

while the expression for pressure is following:

$$p = U_0 \left\{ \omega_0 - \frac{\omega_1}{3} \ln V \right\} V^{(\frac{\omega_1}{6} \ln V - 1)} \exp(-\omega_0 V). \quad (26)$$

The criterion for stability of the fluid during expansion:

1. $\left(\frac{\partial p}{\partial V}\right)_S < 0$.
2. The thermal capacity at constant volume should be greater than zero i.e $c_V > 0$.

The first condition leads to the following constraint:

$$\left(\frac{\omega_1}{3} \ln V - 1 - \omega_0 V\right) \left(\omega_0 - \frac{\omega_1}{3} \ln V\right) < \frac{\omega_1}{3}. \quad (27)$$

Now to get to the thermal equation of state we start with the expression for temperature:

$$T = \left(\frac{\partial U}{\partial S}\right)_V. \quad (28)$$

Using the expression for internal energy we obtain:

$$T = \frac{U}{U_0} \frac{dU_0}{dS} = \frac{dU_0}{dS} V^{(\frac{\omega_1}{6} \ln V)} \exp(-\omega_0 V). \quad (29)$$

Now we will look into U_0 by considering the change of variable from (S,V) to (P,T). The Jacobian of the transformation is (Landau, L. D. and Lifschitz, E. M. 1984):

$$J = \left(\frac{\partial p}{\partial S}\right)_V \left(\frac{\partial T}{\partial V}\right)_S - \left(\frac{\partial p}{\partial V}\right)_S \left(\frac{\partial T}{\partial S}\right)_V. \quad (30)$$

Note, we can also write it out as the following:

$$J = -\left(\frac{\partial T}{\partial S}\right)_V \left(\frac{\partial p}{\partial S}\right)_T. \quad (31)$$

Equating the last two expressions and using the expression for U and T we get the following:

$$\left(\omega_0 - \frac{\omega_1}{3} \ln V\right) \left(-\omega_0 + \frac{\omega_1}{3V} \ln V\right) \left\{ U_0 \frac{d^2 U_0}{dS^2} - \left(\frac{dU_0}{dS} \right)^2 \right\} = 0 \quad (32)$$

$$\Rightarrow U_0 \frac{d^2 U_0}{dS^2} - \left(\frac{dU_0}{dS} \right)^2 = 0 \quad (33)$$

$$\Rightarrow U_0 = \exp(\alpha S), \quad (34)$$

where α is integration constant upon integrating the differential equation for U_0 . Hence we find

$$\rho = \frac{T}{\alpha V} \quad \& \quad (35)$$

$$p = \omega \rho = \frac{T}{\alpha V} \left(\omega_0 - \frac{1}{3} \omega_1 \ln V \right) \quad (36)$$

The existence of critical point requires the following condition to be true:

$$\left. \begin{aligned} \left(\frac{\partial p}{\partial V} \right)_T &= 0 \\ \left(\frac{\partial^2 p}{\partial V^2} \right)_T &= 0 \\ \left(\frac{\partial^3 p}{\partial V^3} \right)_T &< 0 \end{aligned} \right\} \quad (37)$$

But in this model, the point where we have $\left(\frac{\partial p}{\partial V} \right)_T = 0$ has the property that second of above criterion does not hold as

$$\left(\frac{\partial^2 p}{\partial V^2} \right)_T = \frac{T}{\alpha} \frac{\omega''(V)}{V} = \frac{\omega_1}{3V^2} \frac{T}{\alpha V} \neq 0, \quad (38)$$

where prime(t) denotes derivative with respect to volume. Hence, in this model universe does not go through any critical point, even if there is a transition from unstable to stable configuration or vice-versa the transition is smooth.

The specific heat at constant volume comes out to be :

$$c_V = T \left(\frac{\partial S}{\partial T} \right)_V = \frac{1}{\alpha}. \quad (39)$$

Choosing $\alpha > 0$ guarantees the fulfillment of second criterion for stability, i.e., $c_V > 0$. while the velocity of sound is given by:

$$v_s^2 = \omega - \frac{\omega_1}{\omega_1 \ln V - 3 - 3\omega_0 V}. \quad (40)$$

Here also $v_s^2 \leq 1$ imposes constraint on the allowed values of ω_0 and ω_1 .

From the expression for U_0 , the entropy is given by:

$$S = \frac{1}{\alpha} \left[\ln \left(\frac{T}{\alpha} \right) + \omega_0 V - \frac{\omega_1}{6} (\ln V)^2 \right]. \quad (41)$$

3.2 Fluid with $\omega(z) = -1 + \frac{1+z}{3} \frac{A_1 + 2A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2}$

The expression for U in this case becomes:

$$U = U_0 \left(A_0 V + 2A_1 V^{2/3} + A_2 V^{1/3} \right) \exp \left\{ \frac{A_1 \arctan \left(\frac{A_0 V^{1/3} + A_1}{\sqrt{A_0 A_2 - A_1^2}} \right)}{\sqrt{A_0 A_2 - A_1^2}} \right\} \quad (42)$$

where U_0 is integration constant. Therefore energy density comes out to be:

$$\rho = U_0 \left(A_0 + 2A_1 V^{-1/3} + A_2 V^{-2/3} \right) \exp \left\{ \frac{A_1 \arctan \left(\frac{A_0 V^{1/3} + A_1}{\sqrt{A_0 A_2 - A_1^2}} \right)}{\sqrt{A_0 A_2 - A_1^2}} \right\} \quad (43)$$

while the pressure is given by

$$p = \omega(V) U_0 \left(A_0 + 2A_1 V^{-1/3} + A_2 V^{-2/3} \right) \exp \left\{ \frac{A_1 \arctan \left(\frac{A_0 V^{1/3} + A_1}{\sqrt{A_0 A_2 - A_1^2}} \right)}{\sqrt{A_0 A_2 - A_1^2}} \right\} \quad (44)$$

where $\omega(z) = \left(-1 + \frac{1+z}{3} \frac{A_1 + 2A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2} \right)$ and $V = a^3 = (1+z)^{-3}$.

Now we have two criterions for stability of the fluid during expansion :

1. $\left(\frac{\partial p}{\partial V} \right)_S < 0$
2. The thermal capacity at constant volume should be nonnegative i.e $c_V > 0$

The first condition yields:

$$\left(A_0 + 2A_1 V^{-\frac{1}{3}} + A_2 V^{-\frac{2}{3}} \right) \left(\frac{\partial \omega}{\partial V} + \omega(V) \frac{A_1 A_0 V^{-\frac{2}{3}}}{3 \left(A_0 V^{\frac{1}{3}} + A_1 \right)^2 + (A_0 A_2 - A_1^2)} \right) < \frac{2}{3} R \quad (45)$$

where $R = \left(A_1 V^{-\frac{4}{3}} + A_2 V^{-\frac{5}{3}} \right) \omega(V)$.

Now we start with the expression for temperature to obtain thermal EoS:

$$T = \left(\frac{\partial U}{\partial S} \right)_V \quad (46)$$

Using equation(42) we obtain:

$$T = \frac{U}{U_0} \frac{dU_0}{dS} = \frac{dU_0}{dS} (A_0 V + 2A_1 V^{2/3} + A_2 V^{1/3}) \exp \left\{ \frac{A_1 \arctan \left(\frac{A_0 V^{1/3} + A_1}{\sqrt{A_0 A_2 - A_1^2}} \right)}{\sqrt{A_0 A_2 - A_1^2}} \right\} \quad (47)$$

Equating (30) and (31) for this system we have,

$$U_0 \frac{d^2 U_0}{dS^2} - \left(\frac{dU_0}{dS} \right)^2 = 0 \Rightarrow U_0 = \exp(\alpha S)$$

This result resembles with (34). Even we can see the expression for energy density will be similar to (35). Expression for pressure in this case is :

$$p = \omega \rho = \left\{ -1 + \frac{1+z}{3} \frac{A_1 + 2A_2(1+z)}{A_0 + 2A_1(1+z) + A_2(1+z)^2} \right\} \frac{T}{\alpha V} \quad (48)$$

The same conditions (37) applies here too which yields the following conditions needed to be satisfied for having critical point:

$$\omega''(V) = 0 \ \& \ \omega'''(V) < 0 \quad (49)$$

The thermal capacity of the system at constant volume becomes :

$$c_V = T \left(\frac{\partial S}{\partial T} \right)_V = \frac{1}{\alpha} \quad (50)$$

If we choose $\alpha > 0$ it automatically guarantees the fulfillment of second criterion for stability in the same way as previous section i.e $c_V > 0$.

While the velocity of sound is given by:

$$v_s^2 = \omega + \frac{\frac{\partial \omega}{\partial V}}{\frac{\partial(\ln p)}{\partial V}} \quad (51)$$

Now $v_s^2 < 1$ imposes constraint on ω_0 and ω_1 .

From the expression for U_0 , we obtain the expression for entropy which is:

$$S = \frac{1}{\alpha} \left[\ln \left(\frac{T}{\alpha} \right) - \ln \left(A_0 V + 2A_1 V^{2/3} + A_2 V^{1/3} \right) - \left\{ \frac{A_1 \arctan \left(\frac{A_0 V^{1/3} + A_1}{\sqrt{A_0 A_2 - A_1^2}} \right)}{\sqrt{A_0 A_2 - A_1^2}} \right\} \right] \quad (52)$$

4 A General Derivation : Independent of Model

Looking deep into the fact that in last section we have observed U_0 comes out to be $\exp(\alpha S)$ independent of the form of $\omega(z)$, here we are giving a proof without assuming any specific form of $\omega(z)$ which makes the result stronger.

We start with integrating energy conservation equation (23)

$$U = U_0 F(V) \quad (53)$$

where $F(V) = \exp \left\{ -\int \frac{\omega(V)}{V} dV \right\}$. Hence P becomes:

$$P = U_0 \frac{F(V)\omega(V)}{V} \quad (54)$$

Now to get to the thermal equation of state we start with the expression for temperature (28) and using the expression for internal energy we obtain:

$$T = \frac{U}{U_0} \frac{dU_0}{dS} = \frac{dU_0}{dS} F(V) \quad (55)$$

Equating (30) and (31) and using the expression for U and T we get the following:

$$\frac{F(V)F'(V)\omega(V)}{V} \left[U_0 \frac{d^2 U_0}{dS^2} - \left(\frac{dU_0}{dS} \right)^2 \right] = 0 \quad (56)$$

$$\Rightarrow U_0 \frac{d^2 U_0}{dS^2} - \left(\frac{dU_0}{dS} \right)^2 = 0 \Rightarrow U_0 = \exp(\alpha S)$$

where α is integration constant upon integrating the differential equation for U_0 . Immediately, the expression for the energy density becomes $\rho = \frac{T}{\alpha V}$ while the pressure is given by

$$p = \frac{\omega T}{\alpha V} \quad (57)$$

The specific heat at constant volume comes out to be $c_V = T \left(\frac{\partial S}{\partial T} \right)_V = \frac{1}{\alpha}$. and the entropy is given by following expression:

$$S = \frac{1}{\alpha} \left[\ln \left(\frac{T}{\alpha} \right) - \ln F(V) \right] \quad (58)$$

So, T should be $\geq \alpha F(V)$ to ascertain $S \geq 0$.

The third law of thermodynamics demands the entropy to go to 0 as temperature approaches zero. This demand translates into following criterion when we use (58):

$$\lim_{T \rightarrow 0} \frac{F(V)}{T} = \frac{1}{\alpha} = c_V \quad (59)$$

So we can conclude $F(V)$ must tend to 0 as universe cools down to absolute zero. Using this information, from equation (59) we can arrive at the fact that $F(V)$ is actually a measure of change of internal energy to effect a small change in temperature around $T = 0$ point.

Physically Interpreting The General Form of U_0 :

Irrespective of the model we have obtained,

$$U_0 = \exp(\alpha S) \quad (60)$$

In its differential form we can write

$$\frac{dU_0}{U_0} = \alpha dS \quad (61)$$

Now note we have $S = k_B \ln \Omega$ where Ω is the number of micro states corresponding to a macrostate of the system. In its differential form it looks very much like (61)

$$\frac{d\Omega}{\Omega} = \frac{1}{k_B} dS \quad (62)$$

So comparing equation (61) and (62), we interpret U_0 as a measure of number of micro states. Now U_0 is related to energy of universe at present and it is physically very plausible to have energy going in proportional to the number of micro states.

5 Instability : Time scale for Onset of Instabilities

He was Einstein in very known history who has made “the blunder” while producing a static universe which was unstable and not observationally supported! Latter instability regarding cosmological, matter creation has been studied (Saslaw, W. C. 1967). General idea of thermodynamics says that negative specific heat indicates an instability(of course thermal).

Logarithmic Model

Here the expression of c_V illustrates two cases of instability:

(1) $\omega < -1$ along with $\omega_1 + 3\omega(1 + \omega) > 0$: Now it can be easily noted that the former criterion implies the later. Hence we have instability for $\omega < -1$. Taking $\omega_0 = -0.995$ and $\omega_1 = 0.25$ (Efstathiou, G. 1999) we would get that $z = -0.019$ marks the onset of instability. As z goes in the opposite direction of time, negative z indicates towards the future. So $z = -0.019$ can be treated as the timescale for onset of instability. This may suggest that our universe is now in a thermally stable equilibrium but is heading towards an instability where small perturbation can lead to catastrophic change.

(2) $-1 < \omega$ along with $\omega_1 + 3\omega(1 + \omega) < 0$: Now $\omega > 0$ does not satisfy this condition, hence we have stability when $\omega > 0$. So the range is narrowed down to $-1 < \omega < 0$ which in turn implies an interesting scenario that universe starts off with a stable state and it retains stability until ω hits 0. Thereafter it will undergo a stage of thermal instability after which again it will be in a thermally stable state when ω drops below -1 .

In short, depending on the signature of $\omega_1 + 3\omega(1 + \omega)$ the instability may occur at $\omega = -1$ ³ or at $\omega = 0$ for which the corresponding z is -0.019 & 52.517 respectively. However, there are many objects which are at a

³this is the point we do theoretically mark as the phantom barrier, beyond which universe may turn towards different singularities like Big Rip(Caldwell et. al 2003; Nojiri et. al. 2005) etc.

redshift of $z = 52.517$. But we are yet to speculate any kind of impacts of thermal phase transition of universe upon them. So based on current knowledge of explaining observational data we can rule out such a transition at $z = 52.517$. Besides, the model is valid for $z \leq 4$ (Efstathiou, G. 1999). So we can not comment on the second possibility staying within the model.

Note there is observation constraint on ω_0 as follows: $\omega_0 > -1$. So we cant have $\omega < -1$ unless we make z slightly negative i.e we are heading towards instability. Also note ω_1 is assumed to be positive to reconcile with the fact that ω should decrease with time.

The expression for speed of sound (14) within this model also shows instability since with ω going below -1 , the square of the speed becomes less than 0. *This is generally termed as adiabatic instability.* (Bean, R et. al. 2008a, 2008b). Bean, R. et al. (2008a, 2008b) concludes the models with nontrivial effective coupling between dark matter and dark energy can lead to exponential growth of small adiabatic instability which is also characterized by negative sound speed squared. As a result even if universe starts with a uniform fluid the instability will bring upon an exponential growth of small density perturbation. In analogy we also can speculate that it might be possible that at our indicated points small density perturbation will occur opposing fundamental phenomenon like the propagation of sound to take place.

ASSS Model

Within ASSS model, the expression for c_V depicts that the onset of instability happens when ω goes below -2 , which in turn implies the following timescale of onset of instability: Taking allowed values of parameter (Alam, U. et. al. 2004a, 2004b) i.e taking $A_0 = 0, A_1 = -1.169, A_2 = 1.67$ we get that the instability happens to set on at $z = -0.02$. The explanation of this scenario will follow the previous discussion.

6 On the Validity of Thermodynamical Laws

Study of thermodynamical laws with the universe as a thermodynamical system has been done in any literature (Setare, M.R. 2006, 2007a; Setare, M.R., Shafei, S. 2006; Setare, M.R., Vagenas, E. C. 2008; Mazumder, N., Chakraborty, S. 2009, 2010; Bhattacharya, S. Debnath, U. 2011). We will just recall the result relevant for our models.

6.1 Validity on Apparent Horizon

The Friedman metric of a isotropic spatially homogeneous universe is given by:

$$ds^2 = -dt^2 + \frac{a^2}{1 - kr^2} dr^2 + r'^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (63)$$

where $r' = ar$.

From this metric we can easily calculate the radius of the apparent horizon (r'_{AH}) which comes out to be :

$$r'_{AH} = ar_{AH} = \frac{1}{\sqrt{H^2 + \frac{k}{a^2}}} \quad (64)$$

Now, in spatially flat universe, $k = 0$ and thereby we have $r'_{AH} = \frac{1}{H}$.

Following Hawking's idea, the temperature associated with the apparent horizon is:

$$T = \frac{\kappa}{2\pi} \quad (65)$$

where κ is surface gravity of the apparent horizon and given by; $\kappa = \frac{1}{r'_{AH}}$.

Hence, the temperature associated with apparent horizon is $\frac{1}{2\pi r'_{AH}}$.

With $k=0$, using equation (7) we obtain

$$H^2 = \frac{8\pi}{3} \rho \quad (66)$$

To determine the validity of first law thermodynamics we first calculate the energy crossing over this apparent horizon in an infinitesimal dt time which comes out to be (Bousso, R. 2006)

$$-dE_{AH} = \frac{4\pi(P+\rho)}{H^2}dt = -\frac{3}{2}\frac{(1+\omega)}{H(1+z)}dz \quad (67)$$

On the apparent horizon, the entropy is:

$$S_{AH} = \frac{A}{4} = \pi r_{AH}^2 \quad (68)$$

Hence the following holds :

$$T_{AH}dS_{AH} = dr'_{AH} \quad (69)$$

$$dr'_{AH} = -\frac{1}{H^2}dH = -\frac{3}{2}\frac{(1+\omega)}{H(1+z)}dz \quad (70)$$

$$\Rightarrow -dE_{AH} = T_{AH}dS_{AH} \quad (71)$$

Hence the first law of Thermodynamics holds on apparent horizon independent of how we are gonna model $\omega(z)$.

Now, Using Gibbs equation we get (Izquierdo, G. and Pavon D. 2006)

$$TdS_I = dE_I + PdV \quad (72)$$

where S_I , E_I are entropy and enrgy density respectively inside the apparent horizon and E_I is given by

$$E_I = \frac{4}{3}\pi r_{AH}^3 \rho = \frac{1}{2}r'_{AH} \quad (73)$$

and the volume bounded by the apparent horizon is

$$V = \frac{4}{3}\pi r_{AH}^3 \quad (74)$$

Therefore using equation (72),(73) and (74), we get

$$dS_I = \pi r'_{AH}(1+3\omega)dr'_{AH} \quad (75)$$

Rewriting the last equation and using the expression for dS_{AH} , we get

$$\frac{d(S_I + S_{AH})}{dz} = -\frac{9\pi}{2H^2}\frac{(1+\omega)^2}{1+z} \quad (76)$$

$$\Rightarrow \frac{d(S_I + S_{AH})}{da} = \frac{9\pi}{2H^2}(1+\omega)^2(1+z) > 0 \quad (77)$$

Hence holds the Generalized Second Law of Thermodynamics if we consider apparent horizon and the volume bounded by it. So the result holds not only for two types of fluid we have discussed in the paper, but also for every fluid obeying $p = \omega(z)\rho$ which is in quite agreement with (Mazumder, N. et. al. 2010).

6.2 Invalidity of Thermodynamic laws on Event Horizon

The event horizon r_{EH} is defined to be

$$r_{EH} = \int_t^\infty \frac{dt}{a} = \frac{1}{1+z} \int_z^{-1} \frac{-1}{H} dz \quad (78)$$

Hence the event horizon has a hawking's temperature of $\frac{1}{2\pi r_{EH}}$ which yields

$$T_{EH}dS_{EH} = dr_{EH} = \frac{-1}{(1+z)H}dz \quad (79)$$

It follows that:

$$dE_{EH} + T_{EH}dS_{EH} = \frac{dz}{(1+z)H} \left(\frac{3}{2}(1+\omega) - 1 \right) \neq 0 \text{ unless } \omega = -\frac{1}{3} \quad (80)$$

Note $\omega = -\frac{1}{3}$ is the marginal point of Strong Energy Condition which is $3p + \rho \geq 0$.

Employing similar technique we would get, (Xing, L. et. al. 2011)

$$\frac{d(S_I + S_{EH})}{dz} = 2\pi r_{EH}^4 H \frac{dH}{dz} + \pi [3r_{EH}^3 H^2 (1+\omega) + 2r_{EH}] \frac{-1}{(1+z)H} \quad (81)$$

Changing the variable to a we get;

$$\frac{d(S_I + S_{EH})}{da} = 2\pi (r_{EH}^4 H) \left(\frac{dH}{da} \right) + \pi [3r_{EH}^3 H^2 (1+\omega) + 2r_{EH}] \frac{1}{aH} \quad (82)$$

Evidently $\frac{d(S_I + S_{EH})}{da}$ is not necessarily positive for any allowed value of z (as we have $\frac{dH}{da}$ in the expression) which, in turn, implies that the second law of thermodynamics breaks down inside the event horizon in quite agreement with Xing, L. et. al. 2011).

7 Brief Summary

So far we have studied the thermodynamics of universe from two different models. It has been revealed that the first model where $\omega(z) = \omega_0 + \omega_1 \ln(1+z)$ does not go through any critical point. In this model Universe started with a high (fig. 1, 2, 3) energy density, pressure and temperature which falls as z becomes 2 from 0. We imposed restriction on allowed values of ω_0 and ω_1 by demanding $v_s \leq 1$. The thermal capacity at constant volume changes sign as ω becomes smaller than -1 , but this transition is smooth without having critical point. Unlike MCG or GCG here pressure depends both on temperature and volume in thermal EoS. From equation (23) we have shown it is possible to have $c_V > 0$ irrespective of model to ascertain a stable universe.

Thereafter we have generalized the results for a general EoS parameter. We have shown that energy density takes the same form as a function of temperature and volume irrespective of model with a general limiting behavior of thermodynamic parameter (59) to satisfy third law of thermodynamics. We have also interpreted U_0 as a measure of number of micro states and argued for its natural plausibility (61) and (62). Moreover, it is intriguing to note that the thermal EoS (57) looks like ideal gas EoS in some way. Also we have determined the asymptotic behavior of $F(V)$ such that third law of thermodynamics holds. It has been shown that the first and second law of thermodynamics hold on apparent horizon not only for our model but also for any model having $p = \omega(z)\rho$. They fail to be valid on event horizon unless ω takes a specific value which is in perfect agreement with (Mazumder, N. et. al.; Xing, L. et. al. 2011).

One important aspect which have been followed in this paper is a try to speculate the instabilities. Considering, the thermodynamical point at which the heat capacity changes its sign, as a transition point we have followed that the point is nothing but the phantom barrier in the case of Log parametrization. Even there we can find the squared sound speed turns out to be negative. Even for ASSS such point arises for negative z , theoretically indicating to some future point. We have also explored the possibility of the fact that at our indicated points small density perturbation will occur such that the fundamental phenomenon like the propagation of sound does not take place at all.

Acknowledgement :

SP would like to acknowledge a debt of gratitude to Indian Academy of Science for financial support and warm hospitality and IISc, Bnagalore for providing research facilities as the work was done during a time of summer visit. RB thanks ISRO grant “ISRO/RES/2/367/10-11” for providing Research Associate Fellowship. Authors are thankful to Prof Banibrata Mukhopadhyay for fruitful discussions.

References

- Alam, U., Sahni, V., Saini, T. D., and Starobinsky, A. A. :- *Mon. Not. Roy. Astron. Soc.* **344** 1057 (2003). [[arXiv:0303009](#)[astro-ph]].
- Alam, U., Sahni, V., Saini, T. D., Starobinski, A. A. :- *Mon. Not. R. Astron. Soc.* **354** 275 (2004a). [[arXiv:0311364](#)[astro-ph]].
- Alam, U., Sahni, V., Starobinski, A. A. :- *JCAP* **0406** 008 (2004b). [[arXiv:0403687v2](#)[astro-ph]].
- Arkani-Hamed, N., Cheng, H. C., Luty, M. A., Mukohyama, S. :- *JHEP* **0405** 074 (2004). [[arXiv:0312099](#)[hep-th]].
- Armendariz - Picon, C., Mukhanov, V. F. and Steinhardt, P. J. :- *Phys. Rev. Lett.* **85** 4438 (2000). [[arXiv:0004134](#)[astro-ph]].
- Bachall, N.A., Ostriker, J.P., Perlmutter, S., Steinhardt, P.J. :- *Science* **284**, 1481 (1999).
- Bean, R., Flanagan, E.E., Troden, M. :- *New J. Phys.* **10** 033006 (2008a). [[arXiv:0709.1124v2](#)[astro-ph]].
- Bean, R., Flanagan, E.E., Troden, M. :- *Phys. Rev. D* **78** 023009 (2008b). [[arXiv:0709.1128v2](#)[astro-ph]].
- Bhattacharya, S. Debnath, U. :- *Int. J. Theor. Phys.* **50** 525(2011) . [[arXiv:1012.5531](#)[gr-qc]].
- Bousso, R. :- *Phys. Rev. D* **71**, 064024 (2005) [[arXiv : 0412197](#)[hep-th]].
- Caldwell, R. R., Dave, R. and Steinhardt, P. J. :- *Phys. Rev. Lett.* **80** 1582 (1998). [[arXiv:9708069](#)[astro-ph]].
- Caldwell, R. R. :- *Phys. Lett. B* **545** 23(2002). [[arXiv:0105318](#)[astro-ph]].
- Caldwell, R. R., Kaminokowsky, M., Weinberg, N. N. :- *Phys. Rev. Lett.* **91** 071301(2003). [[arXiv:0302506](#)[astro-ph]].
- Cardenas, V. H. and Campo, S. D. :- *Phys. Rev. D* **69** 083508 (2004). [[arXiv:0401031](#)[astro-ph]].
- Chevallier, M., Polarski, D. :- *Int. J. Mod. Phys. D* **10**, 213 (2001). [[arXiv:0009008](#)[gr-qc]].
- Cooray, A. R., Huterer, D. :- *ApJ* **513**, L95 (1999). [[arXiv:9901097](#)[astro-ph]].
- Copeland, E.J., Sami, M., Tsujikawa, S. :- *Int. J. Mod. Phys. D* **15**, 1753 (2006). [[arXiv:0603057](#)[hep-th]].
- Efstathiou, G. :- *Mon. Not. R. Astron. Soc.* **310**, 842 (1999).
- Feng, B., Wang, X. L. and Zhang, X. M. :- *Phys. Lett. B* **607** 35 (2005). [[arXiv:0404224](#)[astro-ph]].
- Ferreira, P. G. and Joyce, M :- *Phys. Rev. D* **58** 023503 (1998). [[arXiv:9711102](#)[astro-ph]].
- Garnavich, P. M. et al :- *Astrophys. J.* **493**, L53 (1998). [[arXiv:9710123](#)[astro-ph]].
- Gong, Y., Wang, B., Wang, A. :- *J. Cosmol. Astropart. Phys.* **01**, 024 (2007). [[arXiv : 0610151](#) [gr-qc]].
- Guo, Z. K., Piao, Y. S., Zhang, X. M. and Zhang, Y. Z. :- *Phys. Lett. B* **608** 177 (2005). [[arXiv:0410654](#)[astro-ph]].
- Izquierdo, G., Pavon D., :- *Phys. Lett. B* **633**, 420 (2006). [[arXiv : 0505601](#) [gr-qc]].
- Jassal, H. K., Bagla, J. S., Padmanabhan, T. :- *MNRAS* **356**, L11 (2005). [[arXiv:0208512](#)[astro-ph]].
- Kamenshchik, A. Y., Moschella, U. and Pasquier, V. :- *Phys. Lett. B* **511** 265 (2001). [[arXiv:0103004](#)[gr-qc]].
- Landau, L. D., Lifschitz, E. M. :- *Statistical Physics third ed., Course of the theoretical physics Vol 5* Butterworth-Heinemann, London, (1984).
- Li, M., Li, X. -D., Wang, S., Wang, Y. :- [[arXiv:1103.5870v4](#)[astro-ph.CO]].
- Linder, E. V. :- *Phys. Rev. Lett.* **90**, 091301(2003) . [[arXiv:0208512](#)[astro-ph]].
- Maor, I., Brustein, R., McMahon, J., Steinhardt, P. J. :- *Phys. Rev. D* **65** 123003(2002). [[arXiv:0112526](#)[astro-ph]].
- Maor, I. and Brustein, R. :- *Phys. Rev. D* **67** 103508 (2003). [[arXiv:0209203](#)[hep-th]].
- Mazumder, N., Biswas, R., Chakraborty, S. :- *Mod. Phys. Lett. A* **25**, 2333 (2010).
- Mazumder, N., Chakraborty, S. :- *Class. Quant. Grav.* **26** 195016 (2009) .
- Mazumder, N., Chakraborty, S. :- *Gen. Rel. Grav.* **42** 813-820 (2010). [[arXiv:1005.3403](#)[gr-qc]].
- Melchiorri, A., Mersini, L. and Trodden, M. :- *Phys. Rev. D* **68**, 043509 (2003). [[arXiv:0211522](#)[astro-ph]].
- Myung, Y. S. :- *Astrophys.Space Sci.* **335**, 561(2011). [[arXiv : 0812.0618v2](#) [gr-qc]].
- Nojiri, S., Odintsov, S. D., Tsujikawa, S. :- *Phys. Rev. D* **71** 063004(2005). [[arXiv : 0501025](#) [hep-th]].
- Padmanabhan, T. :- *Phys. Rept.* **380** 235 (2003). [[arXiv:0212290](#)[hep-th]].
- Padmanabhan, T. :- *Current Science* **88**, 1057(2005). [[arXiv:0411044](#)[astro-ph]].
- Peebles, P. J. E. and Ratra, B. :- *Astrophys. J.* **325** L17 (1988).
- Perlmutter, S. et al :- *Nature* **391** 51 (1998). [[arXiv:9712212](#)[astro-ph]].
- Perlmutter, S. et al. :- [**Supernova Cosmology Project Collaboration**], *ApJ* **517**, 565(1999)[[arXiv:9812133](#)[astro-ph]].

Piazza, F. and Tsujikawa, S. :- *JCAP* **0407** 004 (2004). [*arXiv:0405054*[hep-th]].

Riess, A. G. et al. :- [**Supernova Search Team Collaboration**], *Astron. J.* **116**, 1009(1998)[*arXiv:9805201*[astro-ph]].

Riess, A. G. et al :- *Astrophys. J.* **607**, 665 (2004). [*arXiv:0402512*[astro-ph]].

Sahni, V., Starobinsky, A.A. :- *Int. J. Mod. Phys. D* **9** 373 (2000). [*arXiv:9904398*[astro-ph]].

Sahni, V., Starobinsky, A. A. :- *Int. J. Mod. Phys.* **9** 373 (2003).

Sahni, V., Saini, T. D., Starobinsky, A. A. and Alam, U. :- *JETP Lett.* **77**, 201 (2003). [*arXiv:0201498*[astro-ph]].

Sahni, V., Starobinsky, A. :- *Int. J. Mod. Phys. D* **15**, 2105(2006). [*arXiv:0610026*[astro-ph]].

Saslaw, W. C. :- *MNRAS* **136** 39(1967).

Seljak, U., Slosar, A., McDonald, P.:- *JCAP* **0610**, 014(2006). [*arXiv:0604335*[astro-ph]].

Sen, A. :- *JHEP* **0207** 065 (2002). [*arXiv:0203265*[hep-th]].

Setare, M.R. :- *JCAP* **0609** 011 (2006). [*arXiv:0701242v1*[hep-th]].

Setare, M.R., Shafei, S. :- *Phys.Lett. B* **641** 130 (2006). [*arXiv:0611165v1*[hep-th]].

Setare, M.R. :- *JCAP* **0701** 023 (2007a). [*arXiv:0611165*[hep-th]].

Setare, M., Bouhmadi-Lopez, M.R. :- *Phys.Lett B* **648**, 329 (2007).

Setare, M. R. :- *Eur. Phys. J. C* **52** 689(2007b).

Setare, M.R., Vagenas, E. C. :- *Phys.Lett.B* **666** 111(2008). [*arXiv:0801.4478v2*[hep-th]].

Steinhardt, P. J., Wang, L. and Zlatev, I. :- *Phys. Rev. Lett.*, **59** 123504 (1999). [*arXiv:9812313*[astro-ph]].

Urena-Lopez, L.A., Matos, T. :- *Phys. Rev. D* **62** 081302 (2000). [*arXiv:0003364*[astro-ph]].

Vazqueza, J. A., Bridgesa, M., Hobson, M.P., Lasenby, A.N. :- [*arXiv:1205.0847v1*[astro-ph]].

Xing, L., Chen, J., Gui, Y., Schlegel, E. M., Lu, J. :- *Mod. Phys. Lett. A* **26**, 885 (2011).

Wang, B., Gong, Y., Abdalla, E. :- *Phys. Rev. D* **74**, 083520 (2006). [*arXiv : 0511051*[gr-qc]].

Zlatev, I., Wang, L. and Steinhardt, P. J. :- *Phys. Rev. Lett.* **82** 896 (1999). [*arXiv:9807002*[astro-ph]].